

**A-41**

**Pressure Dependence of Atomic Structure, Magnetism, and Elastic Properties of Fe<sub>3</sub>C: a Combined Synchrotron X-ray Diffraction and Nuclear Resonant Scattering Study**

**Lili Gao<sup>1,2</sup>, Bin Chen<sup>1,3</sup>, Przemyslaw K. Dera<sup>4</sup>, Barbara Lavina<sup>4,5</sup>, Vitali B. Prakapenka<sup>4</sup>, Esen E. Alp<sup>2</sup>, Wolfgang Sturhahn<sup>2</sup>, Jiyong Zhao<sup>2</sup>, and Jie Li<sup>1,6</sup>**

<sup>1</sup>Department of Geology, University of Illinois at Urbana-Champaign, Urbana, IL 61801

<sup>2</sup>Advanced Photon Source, Argonne National Laboratory, Argonne, IL 60439

<sup>3</sup>Now at Seismological Lab, California Institute of Technology, Pasadena, CA 91125

<sup>4</sup>Center for Advanced Radiation Sources, University of Chicago, Chicago, IL 60637

<sup>5</sup>Now at Department of Geoscience, University of Nevada, Las Vegas, NV 89154

<sup>6</sup>Now at Department of Geological Sciences, University of Michigan, Ann Arbor, MI 48109

In this work, we demonstrate the feasibility of studying atomic structure, magnetism, and elastic properties with combined synchrotron x-ray diffraction (XRD) and nuclear resonant scattering methods on Fe<sub>3</sub>C at beamlines 3-ID, 11-BM, 13-BM, and 13-ID at the Advanced Photon Source. XRD probes the long-range order of a material and is a classic method for measuring density. From density-versus-pressure data, elastic properties such as bulk modulus and its pressure derivative can be derived from equation-of-state (EoS) fitting. For EoS fitting, we developed a Java program, FitEoS, which is available to the public. Nuclear resonant scattering technique includes synchrotron Mössbauer spectroscopy and nuclear resonant inelastic x-ray scattering methods. These methods provide information on magnetic properties and lattice vibration. A combination of these methods provides a unique understanding on the behavior of a material under pressure from various perspectives. In this study, we observed two discontinuities in the XRD compression data on Fe<sub>3</sub>C. These two discontinuities were suggested to be related to magnetic and electronic transitions from our nuclear resonant scattering measurements. The implications about the Earth's inner core inferred from this work will also be discussed.